

# Medium-modified fragmentation functions and nPDFs

Rodolfo Sassot  
Universidad de Buenos Aires


Progress in high pT physics at RHIC, Brookhaven, March 2010

# Medium-modified fragmentation functions and nPDFs

Rodolfo Sassot  
Universidad de Buenos Aires

in collaboration with M. Stratmann and P. Zurita  
Phys.Rev.D81 054001 (2010)

Progress in high pT physics at RHIC, Brookhaven, March 2010



# Motivation

→ high pT hadroproduction in pA and AA: RHIC, LHC

# Motivation


- high pT hadroproduction in pA and AA: RHIC, LHC
- study hadronization in different environments

# Motivation

- high pT hadroproduction in pA and AA: RHIC, LHC
- study hadronization in different environments
- factorization & universality in a nucleus?

# Motivation

- high pT hadroproduction in pA and AA: RHIC, LHC
- study hadronization in different environments
- factorization & universality in a nucleus?
- relevant for the extraction of nPDFs



# Phenomenology:

# Phenomenology:

## Early evidence:

SLAC Phys.Rev.Lett. 40, 1624 (1978)

EMC Z.Phys. C52, 1 (1991)

E665 Phys.Rev. D50, 1836 (1994)

# Phenomenology:

## Early evidence:

SLAC Phys.Rev.Lett. 40, 1624 (1978)

EMC Z.Phys. C52, 1 (1991)

E665 Phys.Rev. D50, 1836 (1994)

## Precise SIDIS:

HERMES Nucl. Phys.B 780 1 (2007);

# Phenomenology:

## Early evidence:

SLAC Phys.Rev.Lett. 40, 1624 (1978)  
EMC Z.Phys. C52, 1 (1991)  
E665 Phys.Rev. D50, 1836 (1994)

## Precise SIDIS:

HERMES Nucl. Phys.B 780 1 (2007);


## Precise dAu:

PHENIX Phys.Rev.Lett.98 172302 (2007).  
STAR Phys.Lett.B616, 8 (2005)  
B637, 161 (2006)

# Phenomenology:

Precise SIDIS:

HERMES Nucl. Phys.B 780 1 (2007);





# Phenomenology:

Precise SIDIS:

HERMES Nucl. Phys.B 780 1 (2007);

$$R(z, Q^2, \nu) = \frac{\left( \frac{N^{sidis}}{N^{inc}} \right)_A}{\left( \frac{N^{sidis}}{N^{inc}} \right)_D}$$






# Phenomenology:

PHENIX Phys.Rev.Lett.98  
172302 (2007).

STAR Phys.Lett.B616, 8 (2005)  
B637, 161 (2006)

O.Grebnyuk, Ph.D.Thesis,  
arXiv:0909.3006.




# Phenomenology:

PHENIX Phys.Rev.Lett.98  
172302 (2007).

STAR Phys.Lett.B616, 8 (2005)  
B637, 161 (2006)

O.Grebnyuk, Ph.D.Thesis,  
arXiv:0909.3006.

$$R_{dAu}^H \equiv \frac{1}{\mathcal{N}} \frac{E d^3\sigma^H / dp^3|_{dAu}}{E d^3\sigma^h / dp^3|_{pp}},$$



# Phenomenology:


PHENIX Phys.Rev.Lett.98  
172302 (2007).

STAR Phys.Lett.B616, 8 (2005)  
B637, 161 (2006)

O.Grebnyuk, Ph.D.Thesis,  
arXiv:0909.3006.

$$R_{dAu}^H \equiv \frac{1}{\mathcal{N}} \frac{E d^3\sigma^H / dp^3|_{dAu}}{E d^3\sigma^h / dp^3|_{pp}},$$

$$R_\sigma^H(A, p_T) \equiv \frac{1}{2A} \frac{E d^3\sigma^H / dp^3|_{dA}}{E d^3\sigma^h / dp^3|_{pp}}$$




# Phenomenology:

PHENIX Phys.Rev.Lett.98  
172302 (2007).


STAR Phys.Lett.B616, 8 (2005)  
B637, 161 (2006)

O.Grebnyuk, Ph.D.Thesis,  
arXiv:0909.3006.


$$R_{\sigma}^H(A, p_T) \equiv \frac{1}{2A} \frac{E d^3\sigma^H/dp^3|_{dA}}{E d^3\sigma^h/dp^3|_{pp}}$$




# Phenomenology:




$$R_\sigma^H(A, p_T) \equiv \frac{1}{2A} \frac{E d^3\sigma^H/dp^3|_{dA}}{E d^3\sigma^h/dp^3|_{pp}}$$




# Phenomenology:




$$R_\sigma^H(A, p_T) \equiv \frac{1}{2A} \frac{E d^3\sigma^H/dp^3|_{dA}}{E d^3\sigma^h/dp^3|_{pp}}$$






# Do nuclear effects factorize into PDFs and FFs?




DIS rates to D

$$f_{i/p}(x, Q^2) \longrightarrow f_{i/A}(x, Q^2)$$




Drell Yan rates to D

# Do nuclear effects factorize into PDFs and FFs?




DIS rates to C

$$f_{i/p}(x, Q^2) \longrightarrow f_{i/A}(x, Q^2)$$




scale dependence

# Do nuclear effects factorize into PDFs and FFs?




$$f_{i/p}(x, Q^2) \longrightarrow f_{i/A}(x, Q^2)$$




D.de Florian R.Sassot Phys.Rev.D69 074028 (2004)  
 M.Hirai, S.Kumano, T.H.Nagai Phys.Rev.C76 065207 (2007)  
 K.Eskola, H.Paukkunen, C.A.Salgado, JHEP0904, 065 (2009)

# Do nuclear effects factorize into PDFs and FFs?



DIS rates to C

$$f_{i/p}(x, Q^2) \longrightarrow f_{i/A}(x, Q^2)$$



scale dependence

$$D_{i/p}^h(z, Q^2) \longrightarrow D_{i/A}^h(z, Q^2) ?$$

Do nuclear effects factorize into PDFs and FFs?

Do nuclear effects factorize into PDFs and FFs?

nFFs factorize all non-perturbative details?

Do nuclear effects factorize into PDFs and FFs?

nFFs factorize all non-perturbative details?

“universal” (interchangeable)?

Do nuclear effects factorize into PDFs and FFs?

nFFs factorize all non-perturbative details?

“universal” (interchangeable)?

well defined framework beyond LO?


Do nuclear effects factorize into PDFs and FFs?

nFFs factorize all non-perturbative details?

“universal” (interchangeable)?

well defined framework beyond LO?

constrained by data through global NLO fit?



Why it could not work:

Why it could not work:

factorization breaking

Wiedemann et al. arXiv:1002.2537

non universality of hadronization

Accardi et al. arXiv:0907.3534

modified energy scale dependence

Arleo arXiv:0810.1193

nuclear/high density higher twists


Arleo et al. arXiv:0911.4604

\_\_\_\_\_

\_\_\_\_\_


\_\_\_\_\_

(Please fill in the blanks)



nPDFs digression:


# nPDFs digression:



# nPDFs digression:

## EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)




# nPDFs digression:

EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

nDS nPDFs

D.de Florian R.S.  
Phys.Rev.D69 074028 (2004)



# nPDFs digression:


EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

★ designed to reproduce dAu data

nDS nPDFs

D.de Florian R.S.  
Phys.Rev.D69 074028 (2004)




# nPDFs digression:

EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

★ designed to reproduce dAu data




# nPDFs digression:

## EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

★ designed to reproduce dAu data  
(assuming no FF effects)





# nPDFs digression:

## EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

★ designed to reproduce dAu data  
(assuming no FF effects)

★ unusual gluons





# nPDFs digression:

## EPS nPDFs

K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)

★ designed to reproduce dAu data  
(assuming no FF effects)

★ unusual gluons



# nPDFs digression:

## EPS nPDFs



K.Eskola, H.Paukkunen, C.A.Salgado,  
JHEP0904, 065 (2009)


- ★ designed to reproduce dAu data  
(assuming no FF effects)

- ★ unusual gluons

- ★ extra normalizations: (?)

STAR 0.90  
PHENIX 1.03





Baseline:

consistency  $\rightarrow$  nDS nPDFs

D.de Florian, R.S.

Phys.Rev.D69 074028 (2004)

## Baseline:

consistency  $\rightarrow$  nDS nPDFs

D.de Florian, R.S.  
Phys.Rev.D69 074028 (2004)

reference  $\rightarrow$  DSS FFs

D.de Florian, R.S., M.Stratmann  
Phys.Rev.D75 114010 (2007)  
Phys.Rev.D76 074033 (2007)

# Baseline:


consistency  $\rightarrow$  nDS nPDFs

D.de Florian, R.S.  
Phys.Rev.D69 074028 (2004)

reference  $\rightarrow$  DSS FFs

D.de Florian, R.S., M.Stratmann  
Phys.Rev.D75 114010 (2007)  
Phys.Rev.D76 074033 (2007)

pp  
reference



# Baseline:



consistency  $\rightarrow$  nDS nPDFs

D.de Florian, R.S.  
Phys.Rev.D69 074028 (2004)

reference  $\rightarrow$  DSS FFs

D.de Florian, R.S., M.Stratmann  
Phys.Rev.D75 114010 (2007)  
Phys.Rev.D76 074033 (2007)

pp  
reference



sidis  
reference


# Baseline:

consistency  $\rightarrow$  nDS nPDFs

D.de Florian, R.S.  
Phys.Rev.D69 074028 (2004)

reference  $\rightarrow$  DSS FFs

D.de Florian, R.S., M.Stratmann  
Phys.Rev.D75 114010 (2007)  
Phys.Rev.D76 074033 (2007)




# Baseline:

consistency  $\rightarrow$  nDS nPDFs


D.de Florian, R.S.  
Phys.Rev.D69 074028 (2004)

reference  $\rightarrow$  DSS FFs

D.de Florian, R.S., M.Stratmann  
Phys.Rev.D75 114010 (2007)  
Phys.Rev.D76 074033 (2007)




ATLAS  
arXiv:1003.3124



low pT  
CMS data

CMS  
arXiv:1002.0621



# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

$$W_i(y, A, Q_0^2) = \delta(1 - \epsilon - y)$$

z-shift (energy loss)

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

$$W_i(y, A, Q_0^2) = \delta(1 - \epsilon - y)$$

z-shift (energy loss)

$$W_i(y, A, Q_0^2) = n_i y^{\alpha_i} (1 - y)^{\beta_i}$$

enhancement/suppression, re-shape

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

$$W_i(y, A, Q_0^2) = \delta(1 - \epsilon - y)$$

z-shift (energy loss)

$$W_i(y, A, Q_0^2) = n_i y^{\alpha_i} (1 - y)^{\beta_i}$$

enhancement/suppression, re-shape

weighting coefficients  $\epsilon_i, n_i, \alpha_i, \beta_i$  with a smooth  $A$  dependence

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

$$W_i(y, A, Q_0^2) = \delta(1 - \epsilon - y)$$

z-shift (energy loss)

$$W_i(y, A, Q_0^2) = n_i y^{\alpha_i} (1 - y)^{\beta_i}$$

enhancement/suppression, re-shape

weighting coefficients  $\epsilon_i, n_i, \alpha_i, \beta_i$  with a smooth  $A$  dependence

$$n_i = \lambda^{n_i} + \gamma^{n_i} A^{\delta^{n_i}}$$

with  $\lambda^{n_i}, \gamma^{n_i}, \delta^{n_i}$  parameters to be fitted

# Fitting nFFs: convolution approach

$$D_{i/A}^h(z, Q_0^2) = \int_z^1 dy W_i(y, A, Q_0^2) D_i^h\left(\frac{z}{y}, Q_0^2\right)$$

works for nPDFs  
re-scalings/shifts  
modifies FFs  
natural language NLO

$$W_i(y, A, Q_0^2) = \delta(1 - y)$$

no effects

$$W_i(y, A, Q_0^2) = \delta(1 - \epsilon - y)$$

z-shift (energy loss)

$$W_i(y, A, Q_0^2) = n_i y^{\alpha_i} (1 - y)^{\beta_i}$$


enhancement/suppression, re-shape

weighting coefficients  $\epsilon_i, n_i, \alpha_i, \beta_i$  with a smooth  $A$  dependence

$$n_i = \lambda^{n_i} + \gamma^{n_i} A^{\delta^{n_i}}$$

with  $\lambda^{n_i}, \gamma^{n_i}, \delta^{n_i}$  parameters to be fitted

A very simple example for pion production:



# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2)$$

## Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

$$W_g^\pi(y, A, Q_0^2) = n_g \delta(1 - y) + \epsilon_g \delta(1 - \epsilon_g - y)$$

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

$$W_g^\pi(y, A, Q_0^2) = n_g \delta(1 - y) + \epsilon_g \delta(1 - \epsilon_g - y)$$

$$n_g \neq n_q$$

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

$$W_g^\pi(y, A, Q_0^2) = n_g \delta(1 - y) + \epsilon_g \delta(1 - \epsilon_g - y)$$

$$n_g \neq n_q$$

$$\epsilon_g = \epsilon_q ?$$

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

$$W_g^\pi(y, A, Q_0^2) = n_g \delta(1 - y) + \epsilon_g \delta(1 - \epsilon_g - y)$$

$$n_g \neq n_q$$

$$\epsilon_g = \epsilon_q ?$$

simplest A-dependence:

$$n_q = 1 + \gamma_{n_q} A^{2/3}$$

$$n_g = 1 + \gamma_{n_g} A^{2/3}$$

$$\epsilon_q = \epsilon_g = \gamma_\epsilon A^{2/3}.$$

3 parameters

# Toy parameterization

no significant differences between charged pions

$$W_q^\pi(y, A, Q_0^2) = W_{\bar{q}}^\pi(y, A, Q_0^2) = n_q \delta(1 - y) + \epsilon_q \delta(1 - \epsilon_q - y)$$

z-independent      energy loss?

$$W_g^\pi(y, A, Q_0^2) = n_g \delta(1 - y) + \epsilon_g \delta(1 - \epsilon_g - y)$$

$$n_g \neq n_q$$

$$\epsilon_g = \epsilon_q ?$$

simplest A-dependence:


$$n_q = 1 + \gamma_{n_q} A^{2/3}$$

$$n_g = 1 + \gamma_{n_g} A^{2/3}$$


$$\epsilon_q = \epsilon_g = \gamma_\epsilon A^{2/3}.$$

3 parameters

A	$n_q$	$\epsilon_q = \epsilon_g$	$n_g$
He	0.966	0.001	1.015
Ne	0.902	0.002	1.044
Kr	0.745	0.006	1.115
Xe	0.657	0.008	1.155
Au	0.550	0.010	1.203



# Toy parameterization




— — NLO with nDS nPDFs

— — idem + toy nFFs

# Toy parameterization

normalization & trend




— — NLO with nDS nPDFs

— — idem + toy nFFs

Toy parameterization

normalization & trend




no conflict with  
standard evolution

Toy parameterization


normalization & trend

not flexible enough for  
x-dependence: gluons?

no conflict with  
standard evolution




# Toy parameterization



# Toy parameterization

pT dependence!

quark/gluon interplay





# Toy parameterization

pT dependence!

quark/gluon interplay

$\chi^2/d.o.f. \sim 2$





# Refined parameterization

# Refined parameterization

quark fragmentation

$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1 - y)^{\beta_q} + n'_q \delta(1 - \epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1 - y)^{\beta_g} + n'_g \delta(1 - \epsilon_g - y)$$

gluon fragmentation

# Refined parameterization

quark fragmentation

$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$

gluon fragmentation

$$n_i = \lambda^{n_i} + \gamma^{n_i} A^{\delta^{n_i}}$$

smooth A-dependence

# Refined parameterization

quark fragmentation

$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$

gluon fragmentation

$$n_i = \lambda^{n_i} + \gamma^{n_i} A^{\delta^{n_i}}$$

smooth  $A$ -dependence

$$\begin{aligned}\lambda^n &\sim 1 \\ \lambda^{n'} &\sim 0\end{aligned}$$

vanishing effects as  $A \rightarrow 1$

# Refined parameterization

quark fragmentation

$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$

gluon fragmentation

$$n_i = \lambda^{n_i} + \gamma^{n_i} A^{\delta^{n_i}}$$


smooth  $A$ -dependence

$$\begin{aligned}\lambda^n &\sim 1 \\ \lambda^{n'} &\sim 0\end{aligned}$$

vanishing effects as

$$A \longrightarrow 1$$

$\sim 14$  parameters




$\chi^2 = 396.0$

381 data points

14 parameters

$\chi^2/d.o.f = 1.08$




$\chi^2 = 396.0$

381 data points

14 parameters

$\chi^2/d.o.f = 1.08$

$z$  and  $x$  dependence



$\chi^2 = 396.0$


381 data points

14 parameters

$\chi^2/d.o.f = 1.08$

$z$  and  $x$  dependence

no conflict with  
standard evolution




$\chi^2 = 396.0$

381 data points

14 parameters

$\chi^2/d.o.f = 1.08$




$\chi^2 = 396.0$

381 data points

14 parameters

$\chi^2/d.o.f = 1.08$




$\chi^2 = 396.0$

381 data points

14 parameters


$\chi^2/d.o.f = 1.08$

good description:  
normalization  
pT dependence




$R_{dAu}$  rates are not trivial!

# $R_{dAu}$ rates are not trivial!



# $R_{dAu}$ rates are not trivial!


high-z  
dominated



# $R_{dAu}$ rates are not trivial!

high-z  
dominated

"safe"  
z-range





# $R_{dAu}$ rates are not trivial!

high-z  
dominated

"safe"  
z-range

gluon/quark  
balance






# *z-dependence*

# $z$ -dependence


$$R_q = \frac{D_{q/A}(z, Q^2)}{D_{q/p}(z, Q^2)}$$



# $z$ -dependence

$$R_q = \frac{D_{q/A}(z, Q^2)}{D_{q/p}(z, Q^2)}$$

quarks mimic SIDIS




# $z$ -dependence

$$R_q = \frac{D_{q/A}(z, Q^2)}{D_{q/p}(z, Q^2)}$$

quarks mimic SIDIS

gluons do the opposite




# $z$ -dependence


$$R_q = \frac{D_{q/A}(z, Q^2)}{D_{q/p}(z, Q^2)}$$

quarks mimic SIDIS

gluons do the opposite




low  $z$  behavior not supported by data: artifact?



# A-dependence

# A-dependence




$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1 - \epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1 - \epsilon_g - y)$$

# A-dependence

almost linear




$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1 - \epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1 - \epsilon_g - y)$$

# A-dependence

compensation?

almost linear




$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$

# A-dependence

compensation?

almost linear




$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1 - y)^{\beta_q} + n'_q \delta(1 - \epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1 - y)^{\beta_g} + n'_g \delta(1 - \epsilon_g - y)$$

# A-dependence


compensation?

almost linear



$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$



# Centrality classes

# Centrality classes

nFFs as an average:

# Centrality classes

nFFs as an average:

- ~ underestimate central
- ~ overestimate peripheral

# Centrality classes

nFFs as an average:

- ~ underestimate central
- ~ overestimate peripheral

scale factor

$$c \equiv \left\langle \frac{E d^3 \sigma^{\pi^0} / dp^3 \Big|_{dAu}^{c.c.}}{E d^3 \sigma^{\pi^0} / dp^3 \Big|_{dAu}^{m.b.}} \right\rangle_{c.c.}$$


# Centrality classes


nFFs as an average:

~ underestimate central  
~ overestimate peripheral

scale factor

$$c \equiv \left\langle \frac{E d^3\sigma^{\pi^0}/dp^3|_{dAu}^{c.c.}}{E d^3\sigma^{\pi^0}/dp^3|_{dAu}^{m.b.}} \right\rangle_{c.c.}$$





# Conclusions

# Conclusions

A NLO pQCD factorizable scheme with effective nFF

# Conclusions

A NLO pQCD factorizable scheme with effective nFF  
process-independent universal nFF standard evolution equations

# Conclusions

A NLO pQCD factorizable scheme with effective nFF  
process-independent universal nFF standard evolution equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data

# Conclusions

A NLO pQCD factorizable scheme with effective nFF  
process-independent universal nFF standard evolution equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data  
effective nFFs as tools for “distilling” data

# Conclusions

A NLO pQCD factorizable scheme with effective nFF  
process-independent universal nFF standard evolution equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data  
effective nFFs as tools for “distilling” data  
changes in quark fragmentation “look like” mostly energy loss

# Conclusions

A NLO pQCD **factorizable** scheme with **effective nFF**  
process-independent **universal** nFF **standard evolution** equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data

**effective nFFs** as tools for “distilling” data  
changes in quark fragmentation “look like” mostly energy loss  
effects in gluons are quite different

# Conclusions

A NLO pQCD **factorizable** scheme with **effective nFF**  
process-independent **universal** nFF **standard evolution** equations  
 $\Lambda$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data

**effective nFFs** as tools for “distilling” data  
changes in quark fragmentation “look like” mostly energy loss  
effects in gluons are quite different  
cross sections and rates result from non trivial interplay

# Conclusions

A NLO pQCD factorizable scheme with effective nFF  
process-independent universal nFF standard evolution equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data

effective nFFs as tools for “distilling” data  
changes in quark fragmentation “look like” mostly energy loss  
effects in gluons are quite different  
cross sections and rates result from non trivial interplay

predictions based on nFFs can be tested by upcoming data  
JLAB, RHIC, LHC and in the future at EIC


# Conclusions

A NLO pQCD **factorizable** scheme with **effective nFF**  
process-independent **universal** nFF **standard evolution** equations  
 $A$ ,  $z$ ,  $Q^2$ ,  $v$  and  $pT$ -dependence of SIDIS and dAU data

**effective nFFs** as tools for “distilling” data  
changes in quark fragmentation “look like” mostly energy loss  
effects in gluons are quite different  
cross sections and rates result from non trivial interplay

**predictions based on nFFs** can be tested by upcoming data  
JLAB, RHIC, LHC and in the future at EIC

dAu or pA data can help to further constrain nPDFs,  
serve the reference for AA data,  
provided we have a clear picture of nFF




# Backups

A	$n'_q$	$\epsilon$	$n_q$	$\alpha_q$	$\beta_q$	$n'_g$	$n_g$	$\alpha_g$	$\beta_g$
He	0.949	0.002	0.024	24.183	26.800	1.042	-0.131	19.500	46.241
Ne	0.863	0.006	0.065	23.544	26.131	1.114	-0.351	19.210	41.780
Kr	0.668	0.015	0.157	22.103	24.621	1.276	-0.849	18.557	31.721
Ze	0.564	0.020	0.206	21.331	23.811	1.362	-1.115	18.206	26.331
Au	0.439	0.026	0.265	20.411	22.846	1.466	-1.433	17.789	19.902
$\lambda_i$	1	0	0	24.5607	27.1969	1	0	19.6720	48.8825
$\gamma_i$	-0.0218	0.0010	0.0103	-0.1613	-0.1691	0.0181	-0.0557	-0.0732	-1.1264
$\delta_i$	0.6147	0.6147	0.6147	0.6147	0.6147	0.6147	0.6147	0.6147	0.6147


$$W_q^H(y, A, Q_0^2) = n_q y^{\alpha_q} (1-y)^{\beta_q} + n'_q \delta(1-\epsilon_q - y)$$

$$W_g^H(y, A, Q_0^2) = n_g y^{\alpha_g} (1-y)^{\beta_g} + n'_g \delta(1-\epsilon_g - y)$$



LHC - ATLAS

$\sqrt{s}=900 \text{ GeV}$



# LHC-CMS

